

Supporting Information

Multiparametric Biomechanical and Biochemical Phenotypic Profiling of Single Cancer Cells Using Elasticity Microcytometer

Shuhuan Hu^a, Guangyu Liu^b, Weiqiang Chen^{b,c}, Xiang Li^b, Wei Lu^b, Raymond H. W. Lam^{a,b,d*},
and Jianping Fu^{b,e,f,g*}

^aDepartment of Mechanical and Biomedical Engineering, City University of Hong Kong, Hong Kong; ^bDepartment of Mechanical Engineering, University of Michigan, Ann Arbor, Michigan 48109, USA; ^cDepartment of Mechanical and Aerospace Engineering, New York University, New York 11201, USA; ^dCity University of Hong Kong Shenzhen Research Institute, Shenzhen, China; ^eDepartment of Biomedical Engineering, University of Michigan, Ann Arbor, Michigan 48109, USA; ^fDepartment of Cell and Developmental Biology, University of Michigan, Ann Arbor, Michigan 48109, USA; ^gMichigan Center for Integrative Research in Critical Care, University of Michigan, Ann Arbor, Michigan 48109, USA.

*Correspondence: jpfu@umich.edu, rhwlam@cityu.edu.hk




Figure S1. Left (a) and right (b) confining channel arrays capturing single MCF-10A cells. Most confining channels contained only one cell per channel. In rare cases, multiple single cancer cells were observed to be trapped together in the same confining channels (*e.g.* L10, R23 and R26). These cells were discarded for data analysis. Scale bar, 50 μm .




Figure S2. Histogram of cell diameter for MCF-10A ($n = 180$), MCF-7 ($n = 134$), PC3 ($n = 173$), and HeLa cells ($n = 105$). Data was obtained from three independent experiments.




Figure S3. Representative images showing cell shapes of single MCF-7 (*left*) and PC3 cells (*right*) in confining channels immediately after initial cell trapping (top panel) or 15 min after initial cell trapping (bottom panel). Continuous trapping of MCF-7 and PC3 cells in confining channels coated with anti-EpCAM for 15 min led to some changes of cell shapes with enlarged cell contact areas. Scale bar, 50 μm .




Figure S4. Distribution of single-cell surface EpCAM/anti-EpCAM bond density of the selected cell types ($n > 50$).